

A STUDY OF POSSIBLE TRANSMISSION OF HEPATITIS B VIRUS BY A VECTOR, *Aedes aegypti*: A TRIAL ON RABBIT

Ginsau, M. A and Ahmed, U.A²

Department of Science Laboratory Technology, School of Science and
Technology, Jigawa State Polytechnic, Dutse, Nigeria

² Departments of Biological Science, Sule Lamido University, Kafin Hausa,
Jigawa State, Nigeria

Email: ginsau75@yahoo.com

ABSTRACT

Aedes aegypti were artificially fed with Hepatitis B Virus (HBV) positive blood samples of humans and were allowed to bite rabbits. The rabbits presented signs of infection: geophagy, loss of appetite, thinning, loss of fur and inflammation of liver. This study provides an evidence for transmission of HBV through mosquito. This study is limited to transmission to rabbits, a study of transmission to human is recommended.

Keywords: *Aedes aegypti*, Hepatitis B, Transmission, Virus, Vector

INTRODUCTION

Hepatitis B is the most common cause of viral hepatitis in the world with more than 240million chronic carriers of the virus, 1 million of whom are in the United States (Carroll, 2015). In endemic regions, transmission is thought to be associated with exposure during birth and close contact between young infants (Dienstag, 2015). Risk of infection is highest among intravenous drug users, individuals with high-risk sexual behaviours, healthcare workers, individuals with a history of multiple

transfusions, organ transplant patients, dialysis patients and newborns infected during the birthing process (Carroll, 2015). In approximately two-thirds of patients who develop acute hepatitis B infection, no identifiable exposure is evident (Dienstag, 2015). Close to 780,000 deaths in the world are attributed to hepatitis B. the most endemic regions are in sub-Saharan Africa and East Asia where as much as 10% of adults are chronic carriers (W.H.O., 2014).

Causes of hepatitis can be divided into the following major categories: infectious, metabolic, ischemic, autoimmune, genetic, and other. Infectious agents include viruses, bacteria, and parasites. Toxins, drugs, alcohol, and non-alcoholic fatty liver disease are metabolic causes of liver injury and inflammation. Autoimmune and genetic causes of hepatitis involve genetic predispositions and tend to affect characteristic populations. Parasites can affect liver and activate the immune response, resulting in symptoms of acute hepatitis with increased serum IgE (Harder, 2008). Of the protozoan, *Trypanosoma cruzi*, *Leishmania* species, and the malaria-causing *Plasmodium* species all can cause liver inflammation (Harder, 2008).

Chronic hepatitis presents similarly, but can manifest signs and symptoms specific to liver dysfunction with long-standing inflammation and damage to organ (Khalili, 2013). Acute viral hepatitis follows pattern of infection that involves three distinct phases: 1. *The initial prodromal phase*; 2. *Yellowing of the skin and whites of the eyes* follow the prodrome after about 1 -2 weeks and can last for up to 4 week; and 3. *The recovery phase*,

characterized by resolution of the clinical symptoms of hepatitis (Dienstag, 2014).

Screening for hepatitis is very important as it aimed at identifying people infected with the disease as early as possible (Chou, 2014). Screening consists of a blood test that detects hepatitis B surface antigen (HBsAg). If the (HBsAg) is present, a second test - usually done on the same blood sample (Chou, 2014). For viral hepatitis and other acute causes of hepatitis, the patient's blood test and clinical picture are sufficient for diagnosis (Friedman, 2015).for other causes of hepatitis, especially chronic causes, blood tests may not be useful (Friedman, 2015). In this case, liver biopsy is the gold standard for establishing the diagnosis as histopathologic analysis is able to reveal the precise extent and pattern of inflammation and fibrosis (Friedman, 2015). However, liver biopsy is typically not the initial diagnostic test because it is invasive and is associated with a small but significant risk of bleeding that is increased in patients with liver injury and cirrhosis (Grant, 1999).

Acute hepatitis B infections have a favourable course towards complete recovery in 95- 99% of patients (Dienstag, 2014). However certain factors may portend a poorer outcome, such as co-morbid medical conditions or initial presenting symptoms of ascites, edema, or encephalopathy. Overall, the mortality rate for acute hepatitis is low: ~0.1% in total for cases of hepatitis A and B, but rates can be higher in certain populations (Dienstag, 2014). For prevention of hepatitis B, the Centres for Disease Control (CDC) recommend the routine vaccination of all children under the age of 19 with the hepatitis B vaccine (CDC, 1999).

The first dose starts with the first dose administered as a shot into the muscle before the newborn is discharged from the hospital. An additional two doses should be administered before the child is 18 months (CDC, 2016).

Since mosquitoes can spread some diseases, such as malaria and yellow fever, it is tempting to think they can spread other blood-borne diseases like HIV AIDS and hepatitis (Charles, 2017). In 1974, George and Tzenny have reported on role of mosquitoes in transmission of hepatitis B Virus Infection and their study had involved school children; in 1995, Zheng, *et al.* reported on animal transmission of hepatitis B virus through three genera of mosquitoes and their study involved monkeys, *Tupaia Belangeri*; and in 2015, Hanan had reported that Hepatitis B as Vector-borne transmitted infection but has not ascertain the ability of the insect to be infected as a result of biting of human while we tried transmission of the virus by mosquito to rabbit in this study and this is the first of its kind.

MATERIALS AND METHOD

In an experimental environment, a laboratory- raised species of mosquito, *Aedes aegypti* was artificially fed with human HBV infected blood samples. The mosquitoes were allowed to bite 20 rabbits for one week. The health status of rabbits was instantly subjected to a close physical observation, a month after the bite, their venous blood samples were taken to detect HBsAg and their livers were examined.

RESULTS AND DISCUSSION

A close physical observation of health status of the rabbits revealed geophagy, loss of appetite, thinning, and loss of fur among them. These are suspected to be due to the viral infection and similar to those of human except that they slightly deviate different from what is observed in humans. The symptoms in human involved among others, yellow discoloration of the skin, and poor appetite (*MedilinePlus*, 2016). However, the detection of HBsAg from the serum of the rabbits may prove these symptoms are due to viral infection and considering that HBsAg and anti-HBs were conventional. In the use of test card to test card to check HBsAg, the sensitivity was 88.8% and the specificity was 100% (Fu-yu, *et al.*, 2015). In a different trial, using *Tupaia belangeri*, by Zheng *et al.* (1995), a month after biting, venous blood samples were taken to detect serum HBsA, HBeAg, Anti-HBs and Anti-HBc, among the 29 monkeys, 9 were HBV infective serum markers positive. Moreover, there was evidence that certain species of mosquitoes could transmit hepatitis B virus to animals leading to immunity (Hannan, 2015).

Inflammation of the liver was also observed among the rabbits. This had agreed with what is reported in humans and it is a clear definition of hepatitis as reported by *NIAID* (2016) that hepatitis is inflammation of the liver tissue. Hepatitis is inflammation of the liver tissue and its presentations range from complete asymptomatic to severe liver failure. Among others, the causes include viruses and parasites. In 4 of 16 test monkeys liver biopsies, acute inflammatory changes were found with Haematoxylin-Eosine staining and in the hepatic cells, HBsAg and HBcAg were detected by immunohistochemical double labeling

assays, and HBV-DNA was also positive by in situ hybridization in monkey liver sections (Zheng, 1995). Generally, there is an initial insult that causes liver injury and activation of an inflammatory response, which can become chronic, leading to progressive fibrosis and cirrhosis (Dienstag, 2016). The specific mechanism varies and depends on the underlying cause of the hepatitis. The pathway by which hepatic viruses cause viral hepatitis is best understood in the case of hepatitis B and C (Dienstag, 2016). The viruses do not directly cause apoptosis (cell death). Rather, the infection of liver cells activates the innate and adaptive arms of the immune system leading to an inflammatory response which causes cellular damage and death (Dienstag, 2016).

Moreover, the 20 rabbits put under trial in this study all died within thirty days and this had agreed with what was originally known about the disease as reported by Oon (2012); Trepo (2014) that the ancient Sumerians believed that the liver was the home of the soul, and attributed the findings of jaundice to the attack of one's liver by a devil named Ahhazu. Around 400 B.C., Hippocrates recorded the first documentation of an epidemic jaundice, in particular noting the uniquely fulminant course of a cohort of patients who all died within two weeks. He wrote, "The bile contained in the liver is full of phlegm and blood, and erupts....A such an eruption, the patient soon raves, becomes angry, talks nonsense and barks like a dog".

The result of this study provided an evidence for the possibility of Hepatitis B Virus transmission through mosquito. The question of whether the hepatitis B could be a vector-borne transmitted infection was around in the scientific field since the

1949 (Newkirk, 1975). Towards this, over a period of 29 years a total of 12 researches and two review articles were done to investigate this question. In addition, around five non-English Language articles were published during the same period (Hannan, 2015). The majority of the studies agreed that Hepatitis B Virus could be found in bedbugs for a longer period of time than in the mosquitoes. Moreover, there was evidence that certain species of mosquitoes could transmit Hepatitis B Virus to animals leading to immunity; however, this did not reflect their ability to transmit the virus to humans because it was done under experimental conditions and on a small number of animals (Hannan, 2015).

The role of mosquitoes in transmission of Hepatitis B Virus infection was conducted by George and Tzenny in (1974). They conducted seroepidemiological surveys of school children in Arachova and Anthili, Greece to study the possible role of mosquitoes in transmission of Hepatitis B Virus infection and had reported the overall prevalence of hepatitis B antigen 2.7% in the Anthili population and 0.9% in the Arachova population. And further explained that antibody to Hepatitis B Virus was detected in a significantly higher frequency in the Anthili (20.5%) than in the Arachova (5.1%) population and concluded that this finding gives further support to the previously suggested hypothesis that mosquitoes may play role in the transmission of Hepatitis B Virus infection; In 1975 Newkirk and colleagues studied the fate of HBsAg following a blood meal in two species of mosquitoes (*Aedes aegypti* and *Culex tarsalis*). Both mosquito species were experimentally fed on HBsAg positive blood. In the two mosquito species, the highest

concentrations of HBsAg were detected for a minimum of 20 hours after feeding an infected blood. They suggested that one possible explanation of this was the release of protease trypsin in mosquitoes after blood meal, which destructed the virus. As a result, for the mosquitoes to be effective mechanical transmitters, they would have to be killed during the second feeding within a few hours of complete HBsAg positive blood meal, making this route of transmission of low significance; and in 1995 Zheng *et al.* (1995) conducted study on transmission of Hepatitis B Virus through mosquitoes they fed three genera of mosquitoes with human blood infected with Hepatitis B Virus artificially. The mosquitoes were used to bite monkeys and HBsAg and HBcAg were detected from the hepatic cells of the monkeys and concluded that these results provided an evidence for the possibility of transmission of HBV through mosquitoes and suggest its epidemiological significance in mosquito infested areas.

CONCLUSION

This study proves the vector's role of mosquito in transmission of Hepatitis B Virus infection. This study has also stands an advancement of researches on Hepatitis B Virus transmission by vectors and it filled the wide gap, 1995 - 2018 of research in transmission of Hepatitis B Virus through vector. This study has also presents another dimension of the research, via rabbit.

RECOMMENDATIONS

This study is restricted to transmission to rabbit, advanced researches on transmission to human and zoonotic status are recommended.

REFERENCES

- Bernal, W. and Wendon J. (2013). "Acute Liver Failure". *New England Journal of Medicine*. **369**(26): 2525 - 2534
- Carroll, K. (2015). *Medical Microbiology*. New York, McGraw-Hill. Pp 35
- Centres for Disease Control (1999). "Update: recommendations to prevent hepatitis B virus transmission - United States. MMWR 1999" *cdc.gov preview2016*, **48**:33 - 34
- Centre for Disease Control (2016). "Birth-18 Years and Catchup Immunization Schedules for Providers". *Cdc.gov vaccines*. www.CDC.gov. Retrieved September, 2018
- Charles, D. (2017). Biting Insects Don't Transmit HIV or Hepatitis. www.verywell.com/spreading-hepatitis-by-mosquitoes-17600. Retrieved September, 2018
- Chou, R., Dana, T., Bougatsos, C., Blazina, I., Zakher, B. and Khangura, J. (2014). *Screening for Hepatitis B Virus Infection in Nonpregnant Adolescents and Adults: Systematic Review to Update the 2004 U.S Preventive Services Task Force Recommendation*. www.ncbi.nlm.nih.gov. Retrieved September, 2018
- Dienstag, J.L. (2015). "Acute Viral Hepatitis". *Harrison's Principles of Internal Medicine*, New York, McGraw-Hill, 9th edition Friedman, L. S (2015). *Current Medical Diagnosis & Treatment 2016*. McGraw Hill. Chapter 16

- Fu-yu, W., Yu-Wun, L., Jia-Feng, W., Huey-Ling, C., Hong-Yuan, H., Mei-Hwei, C. and Yen-Hsuan, N. (2016). Pediatrics & Neonatology. **57**(3): 219 -224
- Grant, A. and Neuberger, J. (1999). "Guidelines on the use of liver biopsy in clinical practice" *Gut*, **45** (Suppl 4): 1 - 11
- Harder, A. and Mehlhom H .(2008)." Diseases Caused by Adult Parasites of their District Life CycleStages". *Comparative Hepatitis*. Birkhauser. Pp 161 - 216
- Khalili, M. and Burman, B. (2013). *An Introduction to Clinical Medicine*, McGraw Hill, 7th edition
- MedilinePlus* (2016}. "Hepatitis". [Medilineplus.gov/hepatitis](http://medlineplus.gov/hepatitis). Retrieved September, 2018
- NIAID* (2016)."Hepaptitis".web.archive.org/web/2016. Retrieved September, 2018
- Rutherford , A and Dienstag, J. L. (2016).*CURRENT Diagnosis & Treatment. Gastroenterology, Heptology, & Endoscopy*, New York, McGraw-Hill, third edition
- Oon, G.C. (2017). " Viral hepatitis - the silent killer". *Annals of the Academy of medicine, Singapore*, **41**(7): 279 - 80
- Trepo. C. (2014). " A Brief History of hepatitis milestones". *Liver international*, **34** (Supplement s1): 29 - 37
- World Health Organization (2013). "Hepatitis"
www.who.int/topics/hepatitis/en.2013

World Health Organization (2014). "Hepatitis B"
www.who.int/topics/hepatitis/en. 2014

World Health Organization (2016). "what is Hepatitis?"
www.who.int/topics/hepatitis/en. 2016

Reference to this paper should be made as follows: Ginsau, M. A and Ahmed, U.A² (2019), A Study of Possible Transmission of Hepatitis B Virus by a Vector, *Aedes aegypti*: A Trial on Rabbit. *J. of Biological Science and Bioconservation*, Vol. 11, No. 1, Pp. 43-53
