

## PROXIMATE, PHYTOCHEMICAL QUANTIFICATION AND ANTIDIARRHOEA POTENTIAL OF *GONGRONEMA LATIFOLIUM* EXTRACT IN ALBINO RATS.

<sup>1</sup>Nwachoko, Ndidi., <sup>2</sup>Onuoha, Samuel C., <sup>3</sup>Amadi, Justus O.

<sup>1</sup>Department of Biochemistry, Rivers State University of Science and Technology, Nkpolu-Oroworukwo, Port Harcourt, Rivers State, Nigeria.

<sup>2</sup>Department of Biochemistry, University of Port Harcourt, Choba, Rivers State

<sup>3</sup>Department of chemistry (Biochemistry option), Rivers State University of Science and Technology, Nkpolu-Oroworukwo, Port Harcourt, Rivers State,  
Email: blessedconfidence@yahoo.com

### ABSTRACT

Diarrhoea is a potentially morbid condition with high prevalence worldwide, it constitute a major health situation especially in children under the age of five years old. The world health organization has encouraged studies that can bring about solutions to diarrhoea. Thus this work investigated the proximate, phytochemical quantification and antidiarrhoea potential of *G. latifolium* extract with experimental animals. The result of the proximate analysis reviewed carbohydrate composition to be the highest, followed by the protein composition while fat had the least value. The result of the phytochemical analysis, showed high percentage saponins, flavonoids and alkaloids. Also the result of the antidiarrhoea study showed a reduction in the degree and frequency of defecation of diarrhoea stool. The result indicates that the ethanolic extract of *G. latifolium* contains compounds that could inhibit castor oil induced diarrhoea.

**Keywords:** Diarrhoea, *Gongronema latifolium*, Phytochemicals, Proximate .

### INTRODUCTION

Diarrhoea is a condition of having frequent loosed and watery stool at least three times in a day [1]. It is mostly caused by bacteria

infection. It could also be a symptom of other health cases like in food poisoning. The word diarrhoea is derived from a Greek word dia (through) and rrhoea (flow), meaning flowing through [2]. It is also associated with increase in bowel movement and is characterized by increase in frequency of wet stool sometimes accompanied by stomach cramp [3],[1]. Frequent excretion of formed stools is not diarrhoea but the consistency and sudden release of watery stool denotes diarrhoea condition. Diarrhoea infection is the leading cause of death world wide and continue to take a high public health toll especially in children. In Nigeria, it remains the number one killer disease among children under five (5) years old [4].

*G. latifolium* belongs to the family *Asclepiadaceae*, it is a non wooden, climbing plant with broad green leaf. It has a bitter and slightly sweet taste, used as spice and vegetable in traditional medicine [5]. The plant *Gongronema latifolium* is widely distributed in the tropical rain forest of West African countries such as Nigeria, Ghana, Cameroon, Senegal, Guinea- Bissau, Sierra Leone and Cote d'ivoire. The leaf is used for medicinal and nutritional purposes [6]. In Nigeria, it is popularly called "Utazi" and "Arokeke" by the Igbos and the Yorubas respectively. Various parts of the plant particularly the stem and the leaf are used as chewing sticks or liquor in places like Sierra Leone [7]. Still in Sierra Leone, a decoction of the pounded stem is used for intestinal symptoms that are usually associated with parasitic worms [8].

*G. latifolium* leaf is widely used in folk medicine as spice and vegetable [9]. The leaf is used to prepare food for mothers that have recently put to birth as it is believed that it has the potential to stimulate appetite for food, quickens the return of menstrual cycle and reduce post - partum contraction [10]. Previous studies carried out showed that *G. latifolium* possesses hypoglycemic effect and may be used in managing hyperglycemic conditions and Diabetes mellitus in Nigeria [5]. The presences of certain phytochemical

compounds found in *G. latifolium* suggest that it could be used as a therapeutic plant to combat some health conditions. The leaf is consumed as vegetable, vegetables are edible plants harvested and eaten as supporting food or main dishes and may possess aromatic, bitter or tasteless properties [11]. Eighty percent of some Asian and African countries presently practice the use of herbal medicine for different health issues and this was estimated by the World Health Organization (WHO). Thus this study is aimed at quantifying the phytochemicals present in *G. latifolium* and determination of its antidiarrhoea properties.

## **MATERIALS AND METHODS**

**Materials:** Castor oil (Finest gold) BELL, SONS and CO (DRUGGISTS) LTD, Ethanol (BHD) and Loperamide Hydrochloride. Xepa - Soul Pattinson, Malaysia.

### **Collection of Plant Sample and Preparation**

Fresh green leaf of *Gongronema latifolium* was collected between January and February, 2016 from Rumuosi in Obio/Akpor Local Government Area of Rivers State, Nigeria. The leaf was washed with tap water and spread on a clean flat sack material in a well ventilated room, shade and air dry at room temperature for a week. Dried leaves were ground into fine powdered particles, weighed and stored.

**Proximate Analysis:** [12]. Method was adopted

**Phytochemical Quantification:** [12],[13]. Method was adopted for preparation/extraction of sample for GCms quantitative analysis.

### **Collection of Experimental Animals**

Albino rats weighing between 100- 150g, were obtained from animal house of the University of Port Harcourt, Choba, Nigeria. The animals were acclimatized for a period of seven days prior to the commencement of experiment. The animals were grouped into five (5) different groups of three (3) animals each in each group.

### Castor Oil Induced Diarrhoea in Rats and Faecal Count

Castor oil-induced diarrhoea was determined by the method of [14]. Rats weighing between 100 - 150 g fasted for 18 hours were randomly distributed into five groups. The administration was done. The times taken for onset of diarrhoea and faecal droppings were recorded. Percentage inhibition was calculated [15] - [17].

% inhibition:  $(\text{control} - \text{test}) / \text{control} \times 100$ .

### DISCUSSION

*Gongronema latifolium* is consumed fresh, cooked or dried and applied as powdery spice, it carries a moderate sweet - bitter taste and could be used as bittering agents in brewing to produce characteristic flavor and foam. The leaf could be used as a substitute for commercial hops in lager beer production due to its taste [5]. Table 1 showed the proximate analysis of *Gongronema latifolium*. The result revealed that the plant is a good source of carbohydrate which had the highest nutritional content followed by protein, fibre, moisture and fat had the least value. Earlier studies by [18], reported that *G. latifolium* has high moisture, carbohydrate and protein content.

The quantitative phytochemical analysis of *Gongronema latifolium* presented in Table (2) showed that saponins had the highest value followed by flavonoids, alkaloids, tannin, phenol, oxalate and phytate had the list value. Earlier study suggested that the phytochemical evaluation of the leaf of *G. latifolium* was rich in saponin, alkaloids, phenols and flavonoids [9]. *Gongronema latifolium* is among the type of plants which are good source of potential therapeutic agents that eradicates various diseases due to the presence of bioactive phytochemicals found in the plants [19]. Earlier studies have also reported that the phytochemical compounds like saponin, alkaloids, flavonoids and tannin may be responsible for the antidiarrhoea activity of medicinal plants [20] - [22]. Table (3) showed the inhibitory effect of the ethanolic extract of *G. latifolium* on castor

oil induced diarrhea in experimental animals. The result showed high percentage inhibition of diarrhoea in the groups treated with the extract. The group (3) which received the 200 mg/kg ethanolic extract of the leaf had the highest percentage inhibition followed by group 4 and 5. Comparing the values of the groups pretreated with the plant extract and standard drug both showed inhibitory activity to castor oil induced diarrhoea in the pretreated experimental animals in which the extract especially in the (group 3) exhibited high rate of percentage inhibition of castor oil induced diarrhoea.

It is well documented that castor oil produces diarrhoea due to its most active metabolite - ricinoleic acid by hyper- secretory response, which stimulates peristaltic activity in the small intestine, leading to changes in the electrolyte permeability of the intestinal mucosa [23]. The reduction in faecal output by a substance is the basis of the pharmacological evaluation of a potential antidiarrhoea agent [24],[25]

**Table 1: Proximate Analysis of *Gongronema Latifolium*.**

Nutrient Composition	Percentage (%)
Moisture content	12.49
Ash content	7.36
Fibre content	19.86
Fat content	5.98
Protein content	22.00
Carbohydrate content	32.31

**Table 2: Quantitative phytochemical Analysis of *Gongronema latifolium***

Components	Sub Class	Concentration( $\mu\text{g/g}$ )
Flavonoids	Rutin	5.9438
	Anthocyanin	3.8212
	Catechin	27.78212
	Kaemferol	9.9111
	Epicatechin	1.0695
	(Total) =	48.527
Saponin	Sapogenin	58.2013
Alkaloids	Aspartein	0.0002
	Lunamarine	14.7055
	Ribalinidine	2.3380
	(Total) =	17.04
Tannin		11.6700
Phenol		4.7958
Phytate		0.2948
Oxalate		1.4793

**Table 3: The Effect of Ethanolic Extract of *Gongronema Latifolium* on the Wet Faecal Count of Castor Oil-Induced Diarrhoea in Albino Rats.**

Groups	Treatment	Mean of Wet Faeces	% Inhibition
Groups 1	Untreated	0.0 $\pm$ 0.00	-
Groups 2	Control	2.0 $\pm$ 1.58	-
Groups 3	200mg/kg of extract	0.4 $\pm$ 0.54	81.80
Groups 4	400mg/kg of extract	0.6 $\pm$ 0.56	72.70
Groups 5	5mg/kg/ Loperamide HCl	0.6 $\pm$ 0.56	72.70

## REFERENCES

- WHO (2005). Management of diarrhoea with severe malnutrition, oral and intravenous rehydration.22(14) 33-27
- Ahlquist, D.A. (2001). Constipation and diarrhoea. In; Principles of Internal Medicines. McGraw Hill Medical Publishing Division, New York, pp 241- 247.

- World Health Organization (1999). Diarrhoea disease (online). Retrieved April 30,2011.
- Audu, R.,Umilabug, S.A., Renner, J.K. andAwodiji, J.A. (2000). Diarrhoea management. *Journal of Nigeria Infection Control Association*. 3:15-17.
- Ugochukwu, N.H. and Babady, N.E. (2003). Anti-hyperglycemic effect of aqueous and ethanolic extracts of *Gongronema latifolium* leaves on glucose and glycogen metabolism in livers of normal and streptozotocin- induced diabetic rats. *Life Science*.73(150):1925-1938.
- Dalziel, J.M. (1937). The useful plants of West Tropical Africa. Crown Agents for Colonies, London. p. 230.
- Akuodor, G.C., Idris-Usman, M.S., Mbah, C.C., Megwas, U.A., Akpan, J.L., Ugwu, T.C., Okoroafor, D.O. and Osunkwo, U.A. (2010). Studies in Anti-ulcer, analgesic and anti-pyretic properties of the ethanolic leaf extract of *Gongronema latifolium* Rodients. *African Journal of Biotechnology*. 9(5): 2316 - 2321.
- Deighton, F.C. (1957). Vernacular Botanical Vocabulary for Sierra Leone. Crown Agents for Overseas Government and Administration, London. p. 681.
- Morbise, .O., Fafunso, M.A., Makinde, J.M., Olajide, O.A. and Awe, E.O. (2002). Anti-inflammatory property of the leaves of *Gongronema latifolium*. *Phytotherapy Resource*,16:75 - 77.
- Nwanjo, H.U., Okafor, M.C., and Oze, G.O. (2006).Anti-lipid peroxidative activity of *Gongronema latifolium* instreptozotocin-induced diabetic rats. *Nigeria Journal of Physiological Science*.21(1):61-65.
- Edema, A.O. (1987). Production of Some Common Vegetables. Horticultural Resource Institution Ibadan Nigeria. Pp. 1 - 5.
- AOAC (1990). Association of Official Analytical Chemist.4<sup>th</sup>. Edition.

Washington D.C.

- Harbone, J.B. (1973). *Phytochemical Methods*. Chapman & Hall Ltd, London. pp 188
- Awouters, F, Niemegeers, C.J.E., Lenaerts, F.M. and Jannseen, P.A.J (1978). Delay of castor oil-induced diarrhoea in rats: A new way to evaluate inhibitors of prostaglandins biosynthesis. *Journal of Pharmaceutical Pharmacology*. 30:41-45.
- Izzo A A, Nicholetti M, Giamathasis B, Capasso F (1992). Antidiarrhoea activity of *Ferminalia servicea* Burchex DC Extract in: Capasse, F. and Mascolo, N. (Eds), National Drug and Digestive Tract. EMSI, Rome, 223 - 230
- Mukherjee PK, Saha K, Murugeban T (1998). Screening of Antidiarrhoea Profile of Some Plant Extracts of a Specific Region of West Bengal, India. *Journal of Ethnopharmacology*, 60:85 - 89.
- Karim AH, Maklfia A, Zigyat AL, Bnouham M (2010). Antidiarrhoea activity of crude aqueous extract of *Rubia Tinctorum* L. Roots in rident. *Journal Smooth Muscle Resource*, 46(2): 119 - 123.
- Emebu, P.K. and Anyika, J.U. (2011). Proximate and mineral composition of kale (*Brassica oleracea*) grown in Delta State, Nigeria. *Pakistan Journal of Nutrition*, 10: 190-194.
- Farombi, E.O. (2003). African indigenous plants with chemotherapeutic potentials and biotechnological approach to the production of bioactive prophylactic agents. *African Journal of Biotechnology*. 2(12):66 - 671.
- Galvez. J, Zarzuelo, A, Crespo, M.E, Lorente, M.D., Ocete, M.A, and Jimenez, J. (1993). Anti-diarrhoeal activity of *Euphor biahirta* extract and isolation of an active flavonoid constituent. *Planta Medica*. 59:333-336.



Longanga, O.A., Vercruyssen, A. and Foriers, A. (2000). Contribution to the ethnobotanical, phytochemical and pharmacological studies of

traditionally used medicinal plants in the treatment of dysentery and diarrhoea in Lomela area, Democratic Republic of Congo (DRC). *Journal of Ethnopharmacology*. 60:85-89.

Nwachoko, N. and Jack, I.R. (2015). Phytochemical screening and antidiarrhoea activities of *Tetracarpidium conophrorum* induced in albino rats. *Journal of Biochemistry Research*. Volume 4(4), pp. 021-024.

Bakare, R.I., Magbagbeola, O.A., Akinwande, A.I., Okunowo, O.W. and Green, M. (2011). Antidiarrhoea activity of aqueous leaf extract of *Momordica charantia* in rats. *Journal of Pharmacognosy and Phototherapy*, 3(1): 1-7.

Watson, W. C. and Gordon, R. (1962). Studies on the digestion, absorption and metabolism of castor oil. *Biochemical pharmacology*, 11: 229-236.

Ammon, P.J. and Thomas, P.S. (1974). Effect of oleic and ricinoleic acid net Jajunal water and electrolyte movement. *Journal of Clinical Investigation*, 53:374-379.

---

**Reference** to this paper should be made as follows: <sup>1</sup> Nwachoko, Ndidi., <sup>2</sup> Onuoha, Samuel C., <sup>3</sup> Amadi, Justus O. (2017), Proximate, Phytochemical Quantification and Antidiarrhoea Potential of *Gongronema Latifolium* Extract in Albino Rats. *J. of Biological Science and Bioconservation*, Vol. 9, No. 2, Pp. 25-33

---