
CATCH COMPOSITION OF FIN FISH IN IKERE GORGE, OYO STATE NIGERIA

***Adeosun F.I., I. T. Omoniyi and Y. Akegbejo-Samsons**

Department of Aquaculture & Fisheries Management

University of Agriculture, Abeokuta, Nigeria

Email: adeosunfi@yahoo.com

ABSTRACT

The fin fish composition in Ikere gorge was investigated. The fin fishes comprised a total of 34 species belonging to 13 families. *Chrysichthys nigrodigitatus* was most abundant and highest in biomass (69,198.33kg) accounting for 34.65% of the catch. *Synodontis nigrita*, *Malapterurus electricus*, *Marcusenius psittacus* *Gnathonemus sensgalensis* and *Gnathonemus cyprinoides* were the least in number, while *Phago loricatus* was least in biomass (20.00kg) accounting for 0.01%. The major aim was to assess the fish resources of the gorge and evaluate the water parameters as related to fish production. No significant difference was observed in dissolved oxygen, nitrate, phosphate, water temperature, conductivity, alkalinity, total dissolved solid and pH range value in the stations. The water quality parameters were favourable for fish production. Water level and temperature were observed to guarantee high fish yield in the gorge.

Key word: Fin Fish, Checklist, Ikere gorge.

INTRODUCTION

Aquatic resources are important food and economic resources for many countries. Nigeria is blessed with various aquatic ecosystems, which provide adequate resources for fisheries development system. These ecosystems include rivers, lakes, lagoons and marine environment. Due to unguided management these systems have not been able to sufficiently supply the fish needed to feed the Nigerian populace, which was estimated at about 140million (NPC, 2004). The annual demand for fish in Nigeria is 1.5 million tones. This situation calls for serious and urgent action on how to increase and conserve fish resources to ensure sufficient and sustainable production. Even, at global level fish supply is becoming insufficient as a result of human pressure due to population increases (FAO, 1999). The transition to scarcity of fish cannot be prevented by more intensive fishing but rather could be ameliorated by better management of fisheries resources and interventions to improve equity of resources apart from aquacultural practices. FAO (1999) reported that out of 200,000MT fish stock in all part of the world, more than a quarter is overexploited, depleted or recovering.

Artisanal fisheries in Nigeria represent over 85% of local fish production. In 2004, artisanal sector contributed more than 88% of the domestic fish production out of which artisanal inland water fishery contributed 38.5% (FDF, 2000). At present, there is no biological data for finfish species in Ikere gorge, Oyo State Nigeria. This study paper therefore examines the quantitative and qualitative connection between the catches and the physico-chemical features of Ikere gorge.

MATERIALS AND METHODS

Study Area

The study was carried out in Ikere gorge, 8km east of Ikere village and 30 km North east of Iseyin in Oyo state, Nigeria. The gorge covers all the land between longitude $8^{\circ} 10^1$ and $8^{\circ} 20^1$ E and latitude $3^{\circ}40^1$ and $3^{\circ} 50^1$ N with total area of 300,000km². The monthly rainfall of the gorge is between 0mm-213.7mm per year (Adeosun, 2007). The dry season lasts from November to February with occasional rainfall. The depth and width of the river varies slightly at different sampling stations (Sikoki *et al* 1998). The minimum and maximum widths are 300 and 600meters respectively. Study area was stratified into four zones, using ecological, hydrological strata and intensity of fishing activity as criteria.

Fish Sampling

Data on fish species were collected on monthly basis using multi-fleet gillnet sampling techniques; four sampling stations were randomly selected. This technique involved sampling of each station for the fish composition and abundance using a fleet of eight graded experimental gillnets (mesh sizes from 25.4 to 177.8mm) and Cast net (50.8mm) of similar surface area, which were done simultaneously in the various sampling stations. Samples were taken in the first three days of the first week of every month covering eighteen calendar month of raining and dry season period of the year between July 2004 to December 2005. The first stage involved purposive selection of four hydrological locations in each station with substantial fishing activities.

However, fleets of gill nets were set at dusk and retrieved at dawn in all stations, the investigation were carried out from landings site. The nets were set in different ecological zones in open water, flooded bush patches and shallow bays. In addition, catches from long-lines, fish traps, and hollow cylinders made from bamboo and set hooks were assessed to provide a comprehensive picture of fish species in the lake. The number and weight of fish in each station was measured. The weight of each fish was obtained by weighing samples on a digital mettler balance to the nearest 0.01kg. Sample fishes were counted, sorted out and identified using monographs, descriptions, checklist and keys prepared by Boulenger, (1916), Reed *et al* (1967), Holden and Reed (1972) and Loveque *et al* (1991).

Physico-chemical Parameter

Monthly rainfall data were obtained from the Meteorological Department of Ogun-Osun River Basin Development Authority at Iseyin. Surface water temperature was measured two minutes after dipping mercury- in- glass thermometer at a depth of 5.0cm below the surface water. Depth measurements were made using a graduated rope attached to a lead sinker, lowered from a canoe into the floor of the water. In each month, the average depth of ten pre-determined points were recorded as the water level. Conductivity was measured using a portable meter (Model WTW LF 90), while pH was measured using (Model WTW pH 90) meter. Total dissolved solids (TDS) were measure directly with the Lovibond Tintometer. The dissolved oxygen content was determined using Oxy-Guard Model Mk-11 field oxygen meter. Transparency was measured directly using a Secch disc. Salinity was measured with a Salinometer (Antergo, 28). In taking the salinity, a drop of the test water was placed on the lens of the instrument and allowed to remain for five

minutes. The salinity of the water was then read through the eyepiece. The results of the field experiments was corroborated with the results of laboratory analysis of the physico-chemical parameter.

Results

The results of the physico-chemical characteristics of Ikere gorge between July 2004 and December 2005 are presented in Table 1. There was no significant difference ($p > 0.05$) in the physico-chemical factors of Ikere gorge in the four sampled stations in terms of dissolved oxygen, nitrate, phosphate, temperature, alkalinity and total dissolved solids. However, the mean depth of station 1 was significantly higher ($p < 0.05$) than all the other three stations.

Table 1. The mean values of the Physico-chemical characteristic of Ikere gorge between July 2004 and December 2005.

	Station 1	Station 2	Station 3	Station 4
PH	6.15-8.22	6.08-8.26	6.14-8.62	6.04-8.50
Dissolved Oxygen	7.05±0.22 ^a	6.84±0.22 ^a	6.86±0.23 ^a	6.78±0.30 ^a
Transparency	1.81±0.15 ^a	1.38±0.10 ^b	1.87±0.13 ^a	1.75±0.17 ^{ab}
Nitrate	0.41±0.04 ^a	0.47±0.09 ^a	0.51±0.14 ^a	0.56±0.19 ^a
Phosphate	0.31±0.04 ^a	0.36±0.09 ^a	0.42±0.14 ^a	0.48±0.20 ^a
Temperature	27.33±0.38 ^a	27.44±0.45 ^a	27.55±0.33 ^a	28.04±0.35 ^a
Conductivity	82.86±0.37 ^{ab}	83.39±0.51 ^{ab}	82.49±0.36 ^b	84.24±0.69 ^a
Alkalinity	125.33±6.45 ^a	139.06±11.06 ^a	128.50±5.99 ^a	139.33±12.14 ^a
Total Dissolved Solid	311.88±56.89 ^a	298.00±58.25 ^a	310.71±53.12 ^a	324.83±66.05 ^a
Depth	21.40±2.00 ^a	1.93±0.28 ^b	4.05±0.39 ^b	4.73±0.60 ^b

Table 2 represents the fish species composition in Ikere gorge. A total of 34 species from 13 families were caught during the study period. *Chrysichthys nigrodigitatus* was in highest biomass (69,198kg) accounting for 34.65% of the total fish caught. *Synodontis nigrita*, *Malapterurus electricus*, *Marcusenius psittacus* *Gnathonemus sensgalensis* and *Gnathonemus cyprinoides* were the least in number, while *Phago loricatus* was least in biomass (20.00kg) accounting for 0.01% of the total number of fish caught.

Table 2: Fish species caught and their Percentage Composition of fresh water reaches of Ikere gorge.

Family/Species	No of Fish specimen	% no of specimen	Wt.(kg) of specimen	%Wt. of specimen
BAGRIDAE				
<i>Chrysichthys nigrodigitatus</i>	1423	24.81	69,198.1	34.65
<i>Bagrus docmac niger</i>	46	0.80	1,520.21	0.76

CENTROPOMIDAE				
<i>Lates niloticus</i>	170	2.96	4,312.31	2.16
CHARACIDAE				
<i>Brycinus chaperi</i>	512	8.93	6,290.11	3.15
<i>Brycinus macrolepidotus</i>	30	0.52	2,597.01	1.30
CHANNIDAE				
<i>Parachanna obscura</i>	30	0.52	8,138.22	4.08
CICHLIDAE				
<i>Hemichromis fasciatus</i>	545	9.50	6,170.14	3.09
<i>Sarotherodon galilaeus</i>	432	7.53	20,720.3	10.38
<i>Tilapia melanopleura</i>	596	10.39	26,864.2	13.45
<i>Tilapia zillii</i>	38	0.66	1,449.10	0.73
<i>Tilapia melanotheron</i>	444	7.74	3,893.22	1.95
<i>Tilapia mariae</i>	17	0.30	1,565.14	0.78
<i>Tilapia monody</i>	4	0.06	88.03	0.04
<i>Oreochromis niloticus</i>	33	0.58	7,010.12	3.51
CLARIIDAE				
<i>Clarias gariepinus</i>	39	0.68	1,2611.2	6.32
<i>Heterobranchus bidorsalis</i>	6	0.10	1,254.10	0.63
CYPRINIDAE				
<i>Labeo coubie</i>	12	0.21	414.00	0.21
<i>Barbus occidentalis</i>	368	6.42	3,260.11	1.63
<i>Gara water loti</i>	3	0.05	156.00	0.08
<i>Barilius senegalensis</i>	4	0.07	133.33	0.07
<i>Barilius loati</i>	12	0.21	297.14	0.15
HEPSETIDAE				
<i>Hepsetus odoe</i>	41	0.71	4,777.03	2.39
ICTHYOBORIDAE				
<i>Phago loricatus</i>	2	0.03	20.00	0.01
MOCHOKIDAE				
<i>Synodontis membranaceus</i>	31	0.54	4,061.04	2.03
<i>Synodontis nigrita</i>	1	0.02	47.00	0.02

MALAPTERURIDAE				
Malapterurus electricus	1	0.02	256.00	0.13
MORMYRIDAE				
Mormyrus rume	6	0.10	1,221.01	0.61
Mormyrus deliciosus	23	0.40	544.00	0.27
<i>Hyperopisus bebe occidentalis</i>	21	0.37	4,551.01	0.23
Marcusenius isidori	779	13.58	8,999.23	4.51
Marcusenius psittacus	1	0.02	30.01	0.02
Gnathonemus senegalensis	1	0.02	30.02	0.02
Gnathonemus cyprinoids	1	0.02	55.00	0.03
SCHILBEDAE				
<i>Schilbe mystus</i>	64	1.11	3,463.12	1.73
Total	5736	100	199687.6	100
			5	

The monthly biomass (kg) and number of fish caught during the study period were presented in Table 3. The results show that the peak biomass was recorded in June (23,961kg) accounting for 11.39% of the total monthly fish biomass. The highest number (2,303) accounting for 29.80% of the total number of fish was recorded in February.

Table 3. Monthly Biomass and Number of Fish Caught in Ikere Gorge

Month	Number of Fish	Percentage number of fish	Weight of fish (Kg)	Percentage weight of fish
July 04	276	3.57	11,875	5.66
August	240	3.10	7,187	3.42
September	249	3.22	14,609	6.96
October	301	3.89	10,682	5.09
November	246	3.18	8,339	3.97
December	556	7.19	20,028	9.54
January 05	591	7.65	11,888	5.66
February	2,303	29.80	8,191	3.90

March	215	2.78	9,927	4.73
April	362	4.68	10,159	4.84
May	370	4.79	9,904	4.72
June	476	6.04	23,916	11.39
July	262	3.40	8,976	4.28
August	371	4.80	11,804	5.62
September	174	2.25	10,605	5.05
October	190	2.46	8,970	4.27
November	256	3.31	8,082	3.85
December	301	3.89	14,814	7.06
Total	7,730	100	209,956	100

DISCUSSION

Ikere gorge contained different kinds of fish species. Various sampling stations and fishing gear were used to ensure as comprehensive a sample of ichthyofauna as possible from the study area. From the study a total of 5,736 fish specimens were caught between July 2004 to December 2005. These were identified and classified into 34 species of fish representing 13 families were recorded. This species composition was lower than the results of other studies in other water bodies. Lowe-McConnell, (1964) encountered 44 species on the Rupennine River. Okereke (1990) in her study of Otamiri River Imo State recorded 46 species in 20 families. Other comparable study results include: Sydenham, 1979 of Ogun River (85 species), Reid and Sydenham (1978) of the lower Benue River, Victor and Tetteh (1988) 58 species from Ikpoba River, Imevbore and Okpo (1975) 70 species from River Niger.

The 34 species of fish encountered in this study was however higher than the number recorded by Nweke (1984) in Aba River (29 species). Sydenham (1975) in Odo-ona stream (13 species) and Ekeh (1990) in Nworie River (19 species). The abundance of *Chrysiichthys nigrodigitatus* and commonness of *Tilapia melanopleura* and *Marcuseinus isidori* in this study could be attributed to their ability to tolerate low levels of oxygen and inability to neither bury themselves nor burrow into the muds. The results of the study show Ikere gorge consists of numerous fish species.

The variation in number of species may be explained in three ways: Differences in the physico-chemical conditions cause variations in species composition. Low diversity is a function of low productivity which has been a common feature of small fresh water rivers (Welcomme, 1979). The hypothesis states that high diversity would indicate places of unpredictable hazards or places that would be short lived. Only time would prove how unpredictable nor short lived the lower Num River would be. The difference in the species composition in this study and others may be due to difference in abiotic factors such as temperature, pH, turbidity, dissolved oxygen, conductivity, salinity, alkalinity and nutritive salts. For instance surface water temperature exhibited seasonal variation. pH was neutral to slightly alkaline during the study period. Turbidity was high in the early rainy season and low during the late dry season due to increase in suspended particles in the

river. Dissolved oxygen values are typical of fresh water systems. Conductivity, salinity and alkalinity values are all typical of fresh water systems.

The results from the study show that Ikere gorge consisted of fish species that compared favourably with other fresh water bodies. This information can be used for management decisions and formulation of resource development in the area in addition to the provision of a checklist for fisheries study. The variation in number of species may be explained in two ways: Differences in the physicochemical conditions cause variations in species composition. Low diversity is a function of low productivity which has been a common feature of small fresh water rivers (Welcomme, 1979).

REFERENCES

- Adeosun F.I. (2007). Bio-ecology of the fish fauna of Ikere gorge in Oyo State Nigeria, Ph. D. Thesis, University of Agriculture Abeokuta, Ogun State. 36 pp.
- Boulenger, G. A. (1916). Catalogue of the freshwater fishes of Africa in the British Museum (Nat History). Vols I – IV. London Trustees of the British Museum.
- Ekeh, I.B. (1990). The fish community and water quality of an urban stream – Owerri Southern Nigeria. M.Sc. Thesis. University of Port – Harcourt, 80pp.
- FAO (1999). Management guidelines for Asian flood plain river fisheries. Part 1: A spatial, hierarchical and integrated strategy for adaptive co-management FAO fisheries Technical paper 384/Rome.
- FDF (2000): Memorandum presented by Federal Department of Fisheries (FDF) on the Establishment of Fish Farm Estates in the States and the F.C.T. to the 32nd Regular session of the National fisheries Development Committee (NFDC). Lokoja-Kogi State. 9th-12th Oct.2000.
- Holden, M. and Reed, W. (1972). West African Fresh Water fishes. Longmans Limited. London. 33p.
- Imevbore, A. M. A and W. S. Okpo (1975). Aspects of the biology of Kainji Lake fishes. In: The ecology of Lake Kainji – The transition from river to lake. (Eds: A.M.A. Imevbore and O.S. Adegoke), University of Ife press.
- Loveque, C; Pavgy, O: and Taugels, G.G. (eds) (1991). The fresh and brackish water fishes of West Africa Vol. 1. Musee Royale de. 1 Afriqu centrale. Tervurem. Belgique, Editions de LORSTOM 383pp.
- Lowe-McConnell, R.H. (1964). The fishes of the rupununi Savanna district of British Guiana, South America part 1. Ecological groupings of fish species and effects of seasonal cycle on the fish. *J linn. Sol. Zoo.* 45: 103-144.

- Nweke, E.A. (1984). Fish Community Studies in Aba River M.Sc.Thesis, University of Port Harcourt, Choba. 108pp.
- Okereke, F.O. (1990). Studies on the Fish fauna of Otamiri River Imo State, Nigeria M.Sc.Thesis, University of Port Harcourt, Choba.
- Reed, W. J., A. J. Burchard, J. Hopson and I. Yaro (1967). *Fish and fisheries of Northern Nigeria*. Min. of Agric., Northern Nigeria, 226p.
- Rien, G.M. and Sydenham, H. (1978). A Checklist of lower Benue River fish on Ichthyographical review of the Benue river (West Africa). *J. Nat. Hist.* 13 (1): 41-67.
- Sikoki, F.D.; Hart, A.I. and Abowei, J.F.N. (1998). Gill net Selectivity and fish abundance in Lower Num River, Nigeria. *Journal of Applied Science and Environmental Management* Vol. 1. No. 1 pp 13-19.
- Sydenham, D. H. J. (1975). Observations on the fish populations of a Nigerian forest stream. *Rev. Zool. Afr.* 89: 257-272.
- Welcomme, R.L. (1979). Fishery management in large rivers. FAO Fish Tech. Paper, (194): 60pp.