



---

## ANALYSIS OF DISTURBANCE STORM TIME INDICES

Ahmadu Muhammad Aliyu & Jamaludeen Sambo

Department of Physics

Faculty of Science

Gombe State University, Gombe, Nigeria

Corresponding Author : [hafizmohdz@yahoo.com](mailto:hafizmohdz@yahoo.com)

---

### ABSTRACT

One of the energy sinks during solar wind magnetosphere ionosphere system is ring current injection rate which also depends on the disturbance storm time. In this paper data from world data centre Kyoto obtained at international service for geomagnetic indices was used in this studies and international service for geomagnetism and aeronomy 2002 format was adopted during the analysis. Results obtained shows that the disturbance storm time index varies with time which indicate that proper monitoring of geomagnetic indices is required for accurate energy estimates.

### INTRODUCTION

In the process of energy transfer from the solar wind magnetosphere system, there is a corresponding means of energy dissipation, which include joule heating energy, energy via precipitation and ring current injection that result to magnetic disturbance when interact with earth's magnetic field. The activity of the ring current as a major form of magnetic disturbance is observe by an index known as Disturbance Storm Time (DST) which is a measure in the context of space weather. It gives information about the strength of the ring current around earth caused by solar protons and electrons. The ring current around earth produces a magnetic field that is directly opposite Earth's magnetic field, i.e. if the difference between the solar electrons and protons gets higher, and then Earth's magnetic field becomes weaker.

In an attempt to understand disturbance storm time ,Gannon and Love (2011) produced a 1-min time resolution Storm-Time Disturbance index, the USGS DST, called  $DST^{8507-4SM}$ . The index was based on minute resolution horizontal magnetic field intensity from low-latitude observatories in Honolulu, Kakioka, San Juan and Hermanus, for the years 1985–2007. The method used to produce the index was based on

combination of time and frequency-domain techniques, which more clearly identifies and excises solar quiet variation from the horizontal intensity time series of an individual station than the strictly time domain method used in the Kyoto DST index. The USGS I-min DST was compared against the Kyoto DST, Kyoto Sym-H, and the USGS 1h DST ( $DST^{5807-4SH}$ ). They concluded that, in a time series comparison, Sym-H was found to produce more extreme values and account the variation from the difference in latitude between the observatories. Both Kyoto indices are shown to have a peak in their distributions below zero, while the USGS indices have a peak near zero.

Anh (2005) Provide a method to predict magnetic storm events based on the time series of the DST index over the period 1981–2002. The method is based on the multiple scaling of the measure representation of the DST time series. The measure is modeled as a recurrent iterated function system, which leads to a method to predict storm patterns included in its attractor. Numerical results are provided to evaluate the performance of the method in outside-sample forecasts. Caswell (2014) used a nonlinear autoregressive approach with exogenous input as a novel method for statistical forecasting of the disturbance storm time index, a measure of space weather related to the ring current which surrounds the Earth, and fluctuations in disturbance storm time field strength as a result of incoming solar particles. Given the occurrence of solar activity hours or days before subsequent geomagnetic fluctuations and the potential effects that geomagnetic storms have on terrestrial systems, it would be useful to be able to predict geophysical parameters in advance using both historical disturbance storm time indices and external input of solar winds and the interplanetary magnetic field. By assessing various statistical techniques he determined that artificial neural networks may be ideal for the prediction of disturbance storm time index values which may in turn be used to forecast geomagnetic storms. Unlike the above methods, in this work 2015-2016 hourly provisional data values will be used for the analysis using IAGA-2002 format data obtain at Kyoto, Japan

## MATERIALS AND METHOD

### Materials

#### Data Source

The data used in this research, was obtained from World Data Center (WDC) for Geomagnetism, Kyoto, Japan and converted to IAGA-2002 format by the International Service of Geomagnetic Indices (ISGI) hourly provisional data values were used for 2015 and 2016

### Method

The hourly provisional indices for 2015 and 2016 were converted to mean hourly-daily geomagnetic index as follow to reduce the clusters of data. To minimize error in dealing with a cluster of 1h-cadre of data we centralized them to source K, such that: if  $x_0, x_1, x_2, x_3, \dots, x_{23}$  are the hourly geomagnetic indices for a day, then

$$K = \frac{\sum_{i=0}^{23} X_i}{n} \quad (3.0)$$


Where K is the mean hourly-daily geomagnetic indices and used it to make the analyses.


### Data Analysis

The mean hourly disturbance Storm time obtained from equation 3.0 was used as an input to Microsoft excel spread sheet for analysis. The analysis was carried out first for the year 2015 followed by 2016.

## RESULT AND DISCUSSION

### Results





**Figure 1:** Graphical representation of 2015 and 2016, Disturbance Storm Time

## DISCUSSION

In figure 1, the disturbance happen to be negative at the middle and at the end of 2015, this means that, the DST index are negative, reflecting the westward drift of the energetic, positively charged ions produced during the storm and carrying a westward directed electric current, the same thing happen about 100 days and toward the end of the year (2016). At the 120 days and 270<sup>th</sup> day of 2016, the DST is negative which means it reflects some of its charges. Unlike the case where DST is positive at the 80<sup>th</sup> day of 2015 and 20<sup>th</sup> day of 2016 the charged storm cannot carry westward electric current. But when DST is Zero in both the figures it is indicating the weakness of the earth magnetic field which happen during the early days of 2016, but more frequently during 2015.

## CONCLUSION

This studies show that the Disturbance Storm Time (DST) varies with time, day by day and month by month which clear indicates that the ring current injection as one of the energy sinks varies. This is consistent to many results that the earth magnetic field is not constant.

## RECOMMENDATION

Variation of earth magnetic field is not constant due to the magnetic field of the atmospheric ring current which changes month to month and year to year. Therefore there is need for more research so as to know how to prevent the effect of geomagnetic storm on infrastructure and human's health.

## REFERENCES

- Alabi, O. A. (2015). comparative analysis into the geomagnetic indices (ap, ae, dst, bz) during disturbed period and quiet conditions of solar maximum year (2001). *Journal of space and Scientific Research*, 17., 155-163.
- Anh, V. V. (2005). Nonlinear Processes in Geophysics. *Prediction of magnetic storm events using the Dst index*, 12., 799–806.
- Cash, M. D. (2009). Ring current formation influenced by solar wind substorm conditions. *JOURNAL OF GEOPHYSICAL RESEARCH*, 115., 1-18.
- Caswell, J. m. (2014). A Nonlinear Autoregressive Approach to Statistical Prediction of Disturbance Storm Time Geomagnetic Fluctuations Using Solar Data. *Journal of Signal and Information Processing*, 5., 42-53.
- caswell, J. M. (2014). The nonlinear Autoregressive Approach to statistical Prediction of DST Fluactuation using Solar data. *Journal of Signal and informle process*, 5., 42-53.
- Gannon, J. L. (2011). USGS 1-min Dst index. *tmospheric and solar-terrestrial hysics*, 73., 323-334.
- Gulyaeva, T. L. ( 2014). planetary ionosphere and magnetosphere disturbance storm time events. *Journal of Atmostpheric Physics*, 1-6.
- JoselynD, W. A. (1994). magnetic storm. *journal of solar-terrestrial physics*, 10-16.
- Kivelson, M. R. (1995). Cambridge Univ. press, Cambridge. *Introduction to Space Physics*.
- Kivelson, M.G. and C.T. Russell C.T (1995), *Introduction to Space Physics*, Cambridge Univ. Press, Cambridge

Levitin, A. E. (2016). Physics of Auroral Phenomena. *Physics and mathematical aspect of modeling*, 52-56.